



## King's Research Portal

DOI:

[10.1016/j.jid.2017.06.030](https://doi.org/10.1016/j.jid.2017.06.030)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Tziotzios, C., Ainali, C., Holmes, S., Cunningham, F., Lwin, S. M., Palamaras, I., Bhargava, K., Rymer, J., Stefanato, C., Kirkpatrick, N., Vano-Galvan, S., Petridis, C., Fenton, D. A., Simpson, M. A., Onoufriadis, A., & McGrath, J. A. (2017). Tissue and circulating microRNA co-expression analysis reveals potential involvement of miRNAs in the pathobiology of frontal fibrosing alopecia. *Journal of Investigative Dermatology*, 137(11), 2440-2443. <https://doi.org/10.1016/j.jid.2017.06.030>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Accepted Manuscript

Tissue and circulating microRNA co-expression analysis reveals potential involvement of miRNAs in the pathobiology of frontal fibrosing alopecia

Christos Tziotzios, Chrysanthi Ainali, Susan Holmes, Fiona Cunningham, Su M. Lwin, Ioulis Palamaras, Kapil Bhargava, Janice Rymer, Catherine Stefanato, Niall Kirkpatrick, Sergio Vano-Galvan, Christos Petridis, David A. Fenton, Michael A. Simpson, Alexandros Onoufriadis, John A. McGrath

PII: S0022-202X(17)32736-7

DOI: [10.1016/j.jid.2017.06.030](https://doi.org/10.1016/j.jid.2017.06.030)

Reference: JID 990

To appear in: *The Journal of Investigative Dermatology*

Received Date: 18 April 2017

Revised Date: 27 June 2017

Accepted Date: 27 June 2017

Please cite this article as: Tziotzios C, Ainali C, Holmes S, Cunningham F, Lwin SM, Palamaras I, Bhargava K, Rymer J, Stefanato C, Kirkpatrick N, Vano-Galvan S, Petridis C, Fenton DA, Simpson MA, Onoufriadis A, McGrath JA, Tissue and circulating microRNA co-expression analysis reveals potential involvement of miRNAs in the pathobiology of frontal fibrosing alopecia, *The Journal of Investigative Dermatology* (2017), doi: 10.1016/j.jid.2017.06.030.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Tissue and circulating microRNA co-expression analysis reveals potential involvement of  
miRNAs in the pathobiology of frontal fibrosing alopecia**

Christos Tziotzios<sup>1</sup>, Chrysanthi Ainali<sup>1</sup>, Susan Holmes<sup>2</sup>, Fiona Cunningham<sup>2</sup>, Su M. Lwin<sup>1</sup>,  
Ioulios Palamaras<sup>3</sup>, Kapil Bhargava<sup>1</sup>, Janice Rymer<sup>4</sup>, Catherine Stefanato<sup>5</sup>, Niall Kirkpatrick<sup>6</sup>,  
Sergio Vano-Galvan<sup>7</sup>, Christos Petridis<sup>8</sup>, David A. Fenton<sup>1</sup>, Michael A. Simpson<sup>8</sup>, Alexandros  
Onoufriadis<sup>1</sup>, John A. McGrath<sup>1</sup>

<sup>1</sup>St John's Institute of Dermatology, Division of Genetics and Molecular Medicine, King's  
College London, London, United Kingdom; <sup>2</sup>Alan Lyell Centre for Dermatology, Queen  
Elizabeth University Hospital, Glasgow, United Kingdom; <sup>3</sup>Department of Dermatology,  
Barnet General Hospital, Royal Free Foundation Trust, London, United Kingdom; <sup>4</sup>King's  
College London School of Medicine, King's College London, London, United Kingdom;  
<sup>5</sup>Department of Dermatopathology, St. John's Institute of Dermatology, St. Thomas' Hospital,  
London SE1 7EH, United Kingdom; <sup>6</sup>Department of Plastic Surgery, Guy's Hospital, London  
SE1 9RT, United Kingdom; <sup>7</sup>Ramon y Cajal Hospital, Carretera de Colmenar, KM 9.100,  
Madrid, Spain; <sup>8</sup>Department of Genetics, Division of Genetics and Molecular Medicine,  
King's College London, Guy's Hospital, London SE1 7RT, United Kingdom

**Corresponding author:** John A. McGrath, Dermatology Research Labs, Floor 9 Tower Wing,  
Guy's Hospital, Great Maze Pond, London SE1 9RT, United Kingdom. Tel: 44-2071886409;  
Fax: 44-2071888050; E-mail: [john.mcgrath@kcl.ac.uk](mailto:john.mcgrath@kcl.ac.uk)

**Short title:** microRNAs in frontal fibrosing alopecia

**Abbreviations used:** FFA, frontal fibrosing alopecia; PCA, primary cicatricial alopecia; LPP, lichen planopilaris; eHFSC, epithelial hair follicle stem cells

**Funding sources:** The authors acknowledge financial support from the Department of Health via the National Institute for Health Research (NIHR) Rare Diseases Translational Research Collaboration (NIHR-RD TRC) and a Comprehensive Biomedical Research centre award to Guy's & St Thomas' NHS Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust.

**Conflicts of interest:** None declared.

**Word count:** 998

**Figures:** 2

**Keywords:** Frontal fibrosing alopecia (FFA); miRNA

TO THE EDITOR,

Frontal fibrosing alopecia (FFA) is a predominantly postmenopausal, primary lymphocytic cicatricial alopecia (Kossard, 1994), thought to be a clinical subvariant of lichen planopilaris (Kossard et al., 1997). Somewhat of a misnomer, the descriptor FFA remains widely used for what is a generalized cutaneous lichenoid and fibrosing disorder, invariably associated with widespread body hair loss (Chew et al., 2010), extending beyond the frontovertex, especially if untreated (Figure 1a). Histopathologically, FFA features peri-isthmic lymphocytes (Figure 1b), and immune privilege collapse at the bulge may underpin inflammation-driven epithelial hair follicle stem cell loss and scarring (Tziotzios et al., 2016).

MicroRNAs (miRNAs) are small, non-coding ribonucleic acid molecules (RNAs) that may impact protean biological functions (Lau et al., 2001). They have emerged as potential candidates of pathobiologic, diagnostic and therapeutic interest in chronic inflammatory, fibrotic and autoimmune diseases (Hwang et al., 2006; Galasso et al., 2010; Zhang and Farwell, 2008; Pauley et al., 2009; Patel and Nouredine, 2012). MicroRNAs can be detected in plasma, wherein they circulate in a stable form (Yang et al., 2015; Gilad et al., 2008; Chen et al., 2008; Mitchell et al., 2008; 2008; Valadi et al., 2007; Guo et al., 2017). We sought to characterize miRNAs in FFA and probe disease relevance by undertaking tissue and circulating miRNA co-expression analysis in FFA cases vs matched controls. Neither deep transcriptomic analysis nor miRNA signature exploration has previously been explored for their bearing in FFA, to our knowledge.

Seven newly-diagnosed, treatment-naïve FFA cases and seven matched controls were ascertained for tissue miRNA analysis (Table S1). After written informed consent and institutional ethics approval, temporal scalp skin biopsy was obtained from cases and

controls. Each case was evaluated clinically and histologically to confirm active disease. Part of each biopsy was immediately immersed in RNAlater; total RNA was isolated using RNeasy Plus Universal kit (Qiagen, Valencia, USA) and miRNAs retained according to the manufacturer's protocol (see Supplementary section). All samples underwent microarray analysis on Affymetrix GeneChip miRNA 4.0 arrays. Next, to explore whether circulating miRNAs pertinent to fibrosis are relevant to FFA pathogenesis, venous blood was obtained from a separate cohort of 10 biopsy-proven treatment-naïve FFA cases and 10 matched controls (Table S2). Plasma was isolated by centrifuging blood at 2,000g for 20 minutes at 4°C and stored at -80°C until analysis. MicroRNAs were isolated using Qiagen miRNeasy Serum/Plasma Kit, according to the manufacturer's protocol. Expression analysis was undertaken using the Human Fibrosis miRNA PCR Array (Qiagen) (see Supplementary section).

We first constructed a co-expression network of tissue miRNAs to investigate their molecular importance in FFA. The network was generated by assessing pairwise similarity of miRNAs expressed in each group (controls and FFA), calculated by the Pearson correlation coefficient (PCC) (see Supplementary section and Figure 2a). A co-expression network establishes an association among miRNAs that demonstrate a coordinated expression pattern across a group of samples. It has been postulated that conserved co-expression confers a selective advantage and therefore such genes might be functionally related (Piro et al., 2011; Stuart et al. 2003); such networks can therefore provide insight into how cells accomplish their functions. Only miRNA pairs that correlated above the 0.95 PCC threshold value were represented for each tissue network to capture the strongest relationship for miRNA expression regulation. The resulting FFA network displayed 2,089 co-expressed miRNAs (nodes) with 3,009 non-duplicate interactions (edges), while for controls there were

2,100 co-expressed miRNAs (nodes) with 2,934 non-duplicate interactions (edges). Both tissue networks were used as molecular reference datasets to identify communities of miRNAs sharing common functions. For this analysis, we used the Affinity Propagation Clustering algorithm (see Supplementary section). We generated a list of communities for the two tissue networks and automatically identified cluster centers (exemplars) as representative miRNAs of each community (Table S4).

Next, differential expression analysis was applied in the plasma dataset to calculate the changes in expression between control and FFA samples and to identify up- or down-regulated circulating miRNAs (Figures 2b and S1). Fifty-five miRNAs were found to be up-regulated in FFA while 11 miRNAs were down-regulated. For those circulating miRNAs, we further tested: (i) their predictive value; and (ii) if any of them mapped within the communities of tissue networks. To evaluate their predictive value, we used Random Forest with CARET package in R, which ranks all data elements representing the most influential predictors (Table S5 and Figure S2). To further evaluate circulating miRNAs that were found to be differentially expressed between controls and cases, we mapped them within the tissue-specific co-expression networks. Twenty exemplars in the control tissue miRNA network were significantly enriched in the plasma dataset compared to 27 exemplars, which were significantly enriched in the FFA tissue miRNA network. Amongst these exemplars, there were 17 miRNAs common in both networks (cases and controls): three of these were specific to controls whilst nine were representative of FFA. Of those nine circulating miRNAs, four were found to be highly predictive of disease status. Those miRNAs were, in order of importance, hsa-let-7d-5p, hsa-miR-18a-5p, hsa-miR-20a-5p and hsa-miR-19a-3p, which are also co-targeting similar sets of genes. While the latter three are part of the same cluster, hsa-let-7d-5p is independent of the cluster (Figure 2c) and the commonality in co-targeting

similar genes is of particular interest. As previously suggested, if different miRNAs co-target similar sets of genes, these might regulate the same functions (Tsang et al., 2010; Su et al., 2011). We applied GAGE analysis to infer pathways associated with FFA and found MAPK signaling, Endocytosis and Focal Adhesion pathways to be down-regulated and enriched in the networks of co-targeting genes across these four miRNAs (Table S6).

Our study identified circulating miRNAs hsa-let-7d-5p, hsa-miR-18a-5p, has-miR-20a-5p and hsa-miR-19a-3p as being highly predictive of disease status in FFA by using a skin-specific miRNA network as a validating molecular mapping dataset. Functional validation is needed in future to determine mechanistic relevance, while any additional circulating miRNAs that did not map to the skin network, despite having been found to be predictive in our analysis of plasma, could be further explored in larger cohorts.

### Acknowledgments

The authors acknowledge financial support from the Department of Health via the National Institute for Health Research (NIHR) Rare Diseases Translational Research Collaboration (NIHR-RD TRC) and a Comprehensive Biomedical Research centre award to Guy's & St Thomas' NHS Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust. We are thankful to Clinical Nurse Specialist Ellie Stewart for her contribution to sample collection. We thank our patients and volunteers, as this study would not have been possible without their generous contributions.



## References

- Chen X, Ba Y, Ma L, Cai X, Yin Y, Wang K, et al. Characterization of microRNAs in serum: A novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res* 2008;18:997–1006.
- Chew AL, Bashir SJ, Wain EM, Fenton DA, Stefanato CM. Expanding the spectrum of frontal fibrosing alopecia: a unifying concept. *J Am Acad Dermatol*. 2010;63:653–60.
- Galasso M, Elena Sana M, Volinia S. Non-coding RNAs: a key to future personalized molecular therapy? *Genome Med* 2010;2:12.
- Gilad S, Meiri E, Yogeve Y, Benjamin S, Lebanony D, Yerushalmi Y, et al. Serum microRNAs are promising novel biomarkers *PLoS One* 2008;3:e3148.
- Guo L, Xu K, Yan H, Feng H, Wang T, Chai L, et al. MicroRNA expression signature and the therapeutic effect of the microRNA-21 antagomir in hypertrophic scarring. *Mol Med Rep* 2017;15:1211–21.
- Hwang HW and Mendell JT. MicroRNAs in cell proliferation, cell death, and tumorigenesis. *Br J Cancer* 2006;94:776–80.
- Kossard S. Postmenopausal frontal fibrosing alopecia. Scarring alopecia in a pattern distribution. *Arch Dermatol*. 1994;130:770–4.
- Kossard S, Lee MS, Wilkinson B. Postmenopausal frontal fibrosing alopecia: a frontal variant of lichen planopilaris. *J Am Acad Dermatol*. 1997;36:59–66.
- Lau NC, Lim LP, Weinstein EG, Bartel DP. An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* 2001; 294:858–62.

Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, et al. Circulating microRNAs as stable blood-based markers for cancer detection Proc Natl Acad Sci USA 2008;105:10513–8.

Patel V and Nouredine L. MicroRNAs and fibrosis. Curr Opin Nephrol Hypertens 2012; 21: 410–6.

Pauley KM, Cha S, Chan EK. MicroRNA in autoimmunity and autoimmune diseases. J Autoimmun 2009;32:189–94.

Piro RM, Ala U, Molineris I, Grassi E, Bracco C, Perego GP, Provero P, Di Cunto F: An atlas of tissue-specific conserved coexpression for functional annotation and disease gene prediction. Eur J Hum Genet 2011;19:1173-80.

Stuart JM, Segal E, Koller D, Kim SK. A gene-coexpression network for global discovery of conserved genetic modules. Science 2003;302:249–55.

Su Z, Xia J, Zhao Z. Functional complementation between transcriptional methylation regulation and post-transcriptional microRNA regulation in the human genome. BMC Genomics 2011;12:S5–S15.

Tsang JS, Ebert MS, van Oudenaarden A. Genome-wide dissection of microRNA functions and co-targeting networks using gene set signatures. Mol Cell 2010;38:140–53.

Tziotzios C, Stefanato CM, Fenton DA, Simpson MA, McGrath JA. Frontal fibrosing alopecia: reflections and hypotheses on aetiology and pathogenesis. Exp Dermatol 2016;25:847–52.

Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ, Lotvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 2007;9:654–9.

Yang G, Yang L, Wang W, Wang J, Wang J, Xu Z. Discovery and validation of extracellular/circulating microRNAs during idiopathic pulmonary fibrosis disease progression. *Gene* 2015;562:138–44.

Zhang B and Farwell MA. MicroRNAs: a new emerging class of players for disease diagnostics and gene therapy. *J Cell Mol Med* 2008;12:3–21.

## Figure legends

**Figure 1. Clinical and histologic features of FFA.** (a) Scarring alopecia with frontal and marginal scalp involvement as well as eyebrow loss; (b) Horizontal section of the isthmic area showing perifollicular fibrosis, follicular tufting, and a moderately dense perifollicular lymphoid cell infiltrate. A fragment of a naked hair shaft is also seen. (Hematoxylin-eosin stain; scale bar=75  $\mu$ m). Patient permission to publish these clinical photographs has been received.

**Figure 2. Circulating miRNAs within tissue-specific co-expression network establishes evident differences between FFA and controls.** (a) miRNA expression matrix displays the expression values of the two different phenotypes: FFA (gray color) and Controls (white color) in relation to different miRNAs in tissue-specific dataset. Pairwise similarity in miRNA expression vectors was expressed by the Pearson Correlation Coefficient (PCC). Nodes correspond to miRNAs while edges correspond to the pairwise expression similarity. Tissue co-expression networks in control (left) and FFA (right) were generated. Communities of miRNAs were further detected in the two networks after applying APCluster algorithm. (b) Plasma-specific expression profiles used to identify differentially expressed circulating miRNAs in the skin of patients with FFA, compared to controls. Plasma-specific miRNA signature was further mapped into the cluster centers identified in the two networks after applying APCluster algorithm. Over-expressed miRNAs are colored red while under-expressed miRNAs are depicted in green; (c) Predicted circulating miRNAs co-targeting similar sets of genes.

a



b



